T(I)-Chemistry-H-1B

2021

CHEMISTRY—HONOURS

First Paper

(Group - B)

Full Marks : 50

The figures in the margin indicate full marks. Candidates are required to give their answers in their own words as far as practicable.

CHT-13a

Unit - I

Answer any three questions.

1. (a) The number of gas molecules of mass 'm' with speed between C and C + dC is given by

$$dN = AC^2 e^{-mC^2/2kT} dC$$

Calculate the value of 'A'.

How does the above distribution vary with temperature? $\left(\text{Given}: \int_{0}^{\infty} x^2 e^{-ax^2} dx = \frac{1}{4} \sqrt{\frac{\pi}{a^3}}\right)$

- (b) Calculate the number of binary collision per cm³ per second in helium gas at 27°C and 1 atm. Given that the diameter of He atom = 2×10^{-8} cm. 3+2
- (a) Write down the Maxwell distribution for molecular speed of gases in three dimensions and describe the features using graphical plot.
 - (b) Use the Maxwell's distribution of molecular speed to estimate the fraction of N_2 molecules at 500 K that have speeds in the range of 290–300 ms⁻¹. 3+2
- 3. (a) The virial equation of state in terms of P is given by (neglecting terms of higher order)

$$Z = 1 + \frac{1}{RT} \left(b - \frac{a}{RT} \right) P + \frac{a}{\left(RT \right)^3} \left(2b - \frac{a}{RT} \right) P^2.$$

Set up an expression for the initial slope of 'Z' versus 'P' curve of a real gas and obtain an expression for the Boyle temperature of the gas.

(b) Apply the equipartition principle to calculate $\gamma = \frac{C_P}{C_V}$ for N₂O assuming it to be ideal. 3+2

Please Turn Over

T(I)-Chemistry-H-1B

- 4. (a) Derive the reduced equation of state for 'n' moles of a real gas obeying the van der Waals equation.
 - (b) Consider the following equation of state for a gas :

$$P\overline{V} = RT\left(1 + \frac{b}{\overline{V}}\right)$$

Would it be possible to liquefy the gas? Explain.

- 5. (a) What is Lennard–Jones 6-12 potential? Draw the Lennard–Jones potential against the internuclear distance plot.
 - (b) At normal temperature, iodine is a solid but fluorine is a gas. Explain. 3+2

Unit - II

Answer any two questions.

6. (a) Identify the extensive and intensive properties among the following :

free energy, molar entropy, heat capacity, pressure.

- (b) Calculate Q, W, ΔE and ΔH when one mole of ideal gas is reversibly expanded from 10 atm pressure and 2 lit to 5 atm isothermally at 27°C. 2+3
- 7. (a) Show that $\left(\frac{\partial E}{\partial V}\right)_T = 0$ for an ideal gas.
 - (b) Calculate the enthalpy of formation of NH₃, given the enthalpy of combustion of NH₃ and H₂ are -90.6 kcal mol⁻¹ and -68.3 kcal mol⁻¹ respectively. 2+3
- 8. (a) Derive the integrated form of Kirchhoff's equation to show the variation of ΔH° of a reaction with temperature. Under what condition this variation becomes independent of temperature?
 - (b) The heat capacity of a substance is often given by :

$$\overline{C}_p = a + bT + \frac{C}{T^2}$$

Use this expression to estimate the change in molar enthalpy of carbon dioxide when heated from 20° C to 40° C

Given :
$$a = 44 \cdot 22 \text{ JK}^{-1} \text{ mol}^{-1}$$
, $b = 8 \cdot 79 \times 10^{-3} \text{ JK}^{-2} \text{ mol}^{-1}$, $C = -8 \cdot 02 \times 10^{5} \text{ JK mol}^{-1}$. $3+2$

CHT-13b

Unit - I

Answer any three questions.

- 9. (a) State the two important deductions of Carnot's theorem.
 - (b) Prove both the deductions qualitatively.
- 10. (a) State the two famous statements of second law of thermodynamics by naming them properly.
 - (b) Show that both the statements are equivalent.

2+3

2+3

3+2

2+3

- **11.** (a) What is Clausius Inequality?
 - (b) A certain mass of ideal gas expands into vacuum to twice its initial volume. Calculate ΔU and ΔS of the process. 2+3
- 12. (a) Using Maxwell's relations derive the equation of state : $\left(\frac{\partial H}{\partial P}\right)_T = -T\left(\frac{\partial V}{\partial T}\right)_P + V$
 - (b) Define Joule–Thomson effect and Inversion temperature.
- 13. (a) Differentiate between Joule–Thomson cooling and adiabatic cooling.
 - (b) Draw S-T diagram of a Carnot cycle labelling the diagram, mentioning properly all the processes involved. 2+3

Unit - II

Answer any two questions.

- 14. (a) The half-life period of a first-order reaction is 15 min. Calculate the rate constant and time taken to complete 80% of the reaction.
 - (b) What is a pseudo first-order reaction? Explain with an appropriate example. 2+3
- 15. (a) What is homogeneous catalysis? Explain with a suitable example.
 - (b) Rate constants of a reaction at 300 K and 310 K are 4.5×10^{-5} sec⁻¹ and 9.0×10^{-5} sec⁻¹ respectively. Evaluate the activation energy and pre-exponential factor of the reaction. What should be the order of this reaction? 2+3
- (a) Calculate the activation energy of a reaction, whose rate constant gets tripled by a rise of temperature from 22°C to 32°C.
 - (b) Derive the Michaelis–Menten equation for enzyme catalysis explaining all the terms involved.

2+3