

2021

MATHEMATICS — HONOURS

Paper : DSE-A-1

(Industrial Mathematics)

Full Marks : 65

*The figures in the margin indicate full marks.**Candidates are required to give their answers in their own words as far as practicable.*

1. Choose the correct answer with proper justification / explanation for each of the multiple choice question given below (For each question, one mark for each correct answer and one mark for justification) : 2×10

(a) The attenuation coefficient of an X-ray beam measures

- (i) proportion of the photons absorbed by each millimeter of a substance when an X-ray passes through it.
- (ii) wavelength of the X-ray.
- (iii) proportion of the photons which are not absorbed by a substance when an X-ray passes through it.
- (iv) None of the above.

(b) If $l_{t, \theta}$ be the line through the point $(t \cos \theta, t \sin \theta)$ and perpendicular to the unit vector $\hat{n} = (\cos \theta, \sin \theta)$, then $x + y = \sqrt{2}$ is same as

- (i) $l_{1, \frac{\pi}{2}}$
- (ii) $l_{1, \frac{\pi}{4}}$
- (iii) $l_{0, \frac{\pi}{2}}$
- (iv) $l_{\sqrt{2}, \frac{\pi}{4}}$

(c) Radon transform of $f = e^{-x^2 - y^2}$ is

- (i) $\sqrt{\pi}e^{-p^2}$
- (ii) $\sqrt{\pi}ep^2$
- (iii) πe^{-p^2}
- (iv) $\frac{\sqrt{\pi}}{2}e^{-p^2}$

(d) If f and g be defined and integrable on the real line, the convolution of f and g is defined by

- (i) $(f * g)(x) = \int_{t=-\infty}^{\infty} f(t)g(x+t) dt$ for $x \in \mathbb{R}$
- (ii) $(f * g)(x) = \int_{t=-\infty}^{\infty} f(t)g(x-t) dt$ for $x \in \mathbb{R}$
- (iii) $(f * g)(x) = \int_{t=-\infty}^{\infty} f(t)g(xt) dt$ for $x \in \mathbb{R}$
- (iv) $(f * g)(x) = \int_{t=-\infty}^{\infty} f(t)g(x/t) dt$ for $x \in \mathbb{R}$

Please Turn Over

- (e) Algebraic reconstruction techniques (ARTs) are techniques for reconstructing images
- (i) that have no direct connection to the Radon inversion formula.
 - (ii) that are same as the Radon inversion formula.
 - (iii) that are connected to but not same as the Radon inversion formula.
 - (iv) none of the above
- (f) The Fourier sine transform of $\frac{x}{a^2+x^2}$, a being a constant, is given by
- (i) $2\pi.e^{-ap}$
 - (ii) $\pi^2.e^{-ap}$
 - (iii) $(\pi/2).e^{-ap}$
 - (iv) $\pi.e^{-ap}$
- (g) Let z be a complex number such that $|z| = 4$ and $\arg(z) = 5\pi/6$, then $z =$
- (i) $2\sqrt{3} + 2i$
 - (ii) $-\sqrt{3} + 2i$
 - (iii) $2\sqrt{3} - 2i$
 - (iv) $-2\sqrt{3} + 2i$
- (h) The period of the function $f(x) = \sin(2x) + \frac{1}{\cos(3x)}$ is
- (i) 6π
 - (ii) 2π
 - (iii) π
 - (iv) none of these
- (i) The value of the integral $\int_0^\infty x^5 e^{-x^3} dx$ is
- (i) $1/3$
 - (ii) 1
 - (iii) 0
 - (iv) 2
- (j) If A is a real non-singular symmetric matrix of order n , then
- (i) A and A^{-1} have same set of eigenvectors.
 - (ii) A and A^{-1} have different set of eigenvectors.
 - (iii) A and A^{-1} have some common eigenvectors except one eigenvector.
 - (iv) none of these.

Unit – I

2. Answer **any two** questions :

- (a) What do you mean by X-ray Computerized Tomography (CT)? Explain with example.
- (b) (i) If z is a complex number, then find the minimum value of $|z| + |z - 1|$.

(ii) For any two complex numbers z_1 and z_2 and any real numbers a and b ; then show that

$$|(az_1 - bz_2)|^2 + |(bz_1 - az_2)|^2 = (a^2 + b^2)(|z_1|^2 + |z_2|^2) \quad 2+3$$

(c) Define (in the Hadamard sense) the well-posedness of a mathematical problem. Give an example of an ill-posed problem. 3+2

(d) Solve the following differential equation : $x \frac{d^2 y}{dx^2} + (x-1) \frac{dy}{dx} - y = x^2$. 5

Unit – II

3. Answer **any two** questions :

(a) Direct problem is given by : a continuous function $x : [0, 1] \rightarrow R$, compute $y(t) := \int_0^t x(s) ds$, $t \in [0, 1]$.

Find its inverse problem. 5

(b) Find the inverse function of the function defined by $f(x) = -x^5$, for $x \in [-1, 1]$. Is the inverse function f^{-1} is continuous at $x = 0$? 2+3

(c) Consider the boundary value problem $\frac{d^2 u}{dx^2} = f$, $u(0) = u(l) = 0$, where $f : R \rightarrow R$ is a given continuous function. Suppose that the solution u and f are known. Find the length l of the interval. 5

(d) If 5, 2, 2 are eigenvalues of a square matrix A of order 3 having eigenvectors $a \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$ and

$b \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix} + c \begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix}$ associated with 5 and 2 respectively, where $a \neq 0$, $(b, c) \neq (0, 0)$, then find the matrix A . 5

Unit – III

4. Answer **any one** question :

5×1

(a) An X-ray beam $A(x)$ propagates in a uniform medium which is defined by $A(x) = x$. Prove that the intensity $I(x)$ of this beam is a Gaussian distribution, with boundary conditions $\lim_{|x| \rightarrow \infty} I(x) = 0$. Find the average value of this intensity.

Please Turn Over

- (b) If the intensity of an X-ray light beam is $I(x) = (2x+3)e^{-\frac{dx^2}{2}}$, $x > 0$, then find the inhomogeneous medium and hence show that it is zero at the point $x = -\frac{3}{4} + \frac{\sqrt{16+9d}}{4\sqrt{d}}$, where d is a positive real constant.

Unit – IV

5. Answer **any one** question : 5×1

(a) Show that Radon transform is a linear transform.

- (b) Prove that the line $\mathcal{L}_{1/2, \pi/6}$ has a standard form $x = \frac{\sqrt{3}}{4} - \frac{s}{2}$, $y = \frac{1}{4} + \frac{\sqrt{3}}{2}s$, then find the Random

transformation of $f(x, y) = \begin{cases} x, & x^2 + y^2 \leq 1 \\ 0, & \text{elsewhere} \end{cases}$ at the point $(1/2, \pi/6)$.

Unit – V

6. Answer **any one** question : 5×1

(a) What are the differences between back projection and Random transformation?

(b) Find back projection of the Radon transform of a attenuation-coefficient function f .

Unit – VI

7. Answer **any two** questions : 5×2

(a) Write a short note on algebraic reconstruction technique on the base of CT scan.

(b) If $f(x)$ is an absolutely integrable and piecewise continuous function with a point of discontinuity at $x = \alpha$ but $\lim_{x \rightarrow \alpha^-} f(x)$ and $\lim_{x \rightarrow \alpha^+} f(x)$ exist, then prove that

$$\mathcal{F}^{-1}(\mathcal{F} f)(\alpha) = \frac{1}{2} \left(\lim_{x \rightarrow \alpha^-} f(x) + \lim_{x \rightarrow \alpha^+} f(x) \right).$$

(c) Show that the inverse Fourier transform of an even function is a real-valued function and the inverse Fourier transform of an odd function is a purely imaginary function.

(d) Find the inverse Fourier transform of the function $F(w) = \frac{2}{1+w^2}$.
